

Extinctions and Biodiversity in the Fossil Record

Ricard V Solé and Mark Newman

Volume 2, **The Earth system: biological and ecological dimensions of global environmental change**, pp 297–301

Edited by

Professor Harold A Mooney and Dr Josep G Canadell

in

Encyclopedia of Global Environmental Change
(ISBN 0-471-97796-9)

Editor-in-Chief

Ted Munn

© John Wiley & Sons, Ltd, Chichester, 2002

Extinctions and Biodiversity in the Fossil Record

Ricard V Solé and Mark Newman
Santa Fe Institute, Santa Fe, NM, USA

Life has existed on Earth for more than three billion years. Until the Cambrian explosion about 540 million years ago, however, life was restricted mostly to single-celled microorganisms that were, for the most part, poorly preserved in the fossil record. From the Cambrian explosion onwards, by contrast, we have a substantial fossil record of life's development, which shows a number of clear patterns, including a steady increase in biodiversity towards the present, punctuated by a number of large extinction events that wiped out a significant fraction of the species on the planet, and in some cases, caused major reorganizations amongst the dominant groups of organisms in the ecosphere.

The history of life on the Earth begins in the oceans during the *Achaean eon*, somewhere around 3.8 billion years ago, probably with the appearance first of self-reproducing RNA molecules and subsequently of prokaryotic single-celled organisms. At the start of the *Proterozoic eon*, around 2.5 billion years ago, the atmosphere changed from reducing to oxidizing, as a result of the depletion of stocks of elemental iron in the Earth's crust, and oxygen-breathing life became possible. Eukaryotes, multicellular life, and sexual reproduction all appeared for the first time during the Proterozoic, although the exact order and dates are still in dispute, since the fossil record of this period is poor. The earliest firm evidence of multicellularity dates from about 575 million years ago.

About 540 million years ago, for unknown reasons, an enormous increase in the diversity of multicellular animals took place. This event, known as the *Cambrian explosion*, produced all the major body plans of animals seen today, as well as a number of others that have since become extinct. The first land-dwelling plants appeared during the *Silurian period*, about 430 million years ago, followed shortly afterwards by the first land-dwelling animals, which were insects, and then the first land-dwelling vertebrates during the *Devonian*. About 250 million years ago, at the end of the *Permian* period, the largest mass extinction of all time took place, killing at least 90% of all species on the Earth, and ending the eon named the *Paleozoic*.

The *Mesozoic*, colloquially known as *the age of the dinosaurs*, followed the Paleozoic. In addition to dinosaurs, the Mesozoic also saw the first appearance of mammals and of flowering plants. It ended about 65 million years ago with the *Cretaceous–Tertiary (KT) extinction event*, which

wiped out the dinosaurs along with about 70% of all other species then alive. The interval from the KT event until the present, known as the *Cenozoic eon*, saw the radiation of the mammals to fill many of the dominant land-dwelling niches and, eventually, the evolution of mankind.

This information comes from geological studies and from the substantial fossil record of extinct lifeforms. The currently known fossil record includes about one-quarter of a million species, mostly dating from the time interval between the Cambrian explosion and the present. There are numerous biases in the fossil record, which make accurate quantitative investigations difficult, including the following.

1. Older fossils are harder to find because they are typically buried in deeper rocks than more recent ones.
2. Accurate dating of fossils is difficult. Radiocarbon dating, for example, is not useful for rocks that are hundreds of millions of years old. Radioactive isotopes other than carbon with longer half-lives are used for most of the geologic time-scale, but resolution of dates using these isotopes can be poor.
3. Particularly rich fossil beds, or particularly zealous investigators, may produce very complete records for some time periods or groups of organisms, while other periods or groups may be comparatively poorly researched.
4. Prehistoric environmental disturbances can upset the deposition processes by which fossils are formed and give rise to time periods in which the fossil record is poor.
5. Marine organisms tend to be much better preserved than land-dwelling ones, because deposition is much more uniform and reliable in the oceans than it is on land.

Despite these biases, a number of trends are clear from the fossil record.

Figure 1 shows a plot of the best estimate of the number of living families of marine organisms, as a function of time since the start of the Cambrian. As the plot shows, diversity appears to increase substantially over time, and this is believed to be a real trend. On average, there have been more families, and indeed more species, alive in recent times than earlier. Over the course of the plot, diversity appears to increase by a factor of about five, although some of this increase is an artifact of the greater availability of fossils of recent species in more easily accessible rocks.

Figure 2 shows the extinctions, again of marine organisms, as a function of time over the same interval. The graph gives the number of families becoming extinct per stratigraphic stage. Stages are uneven intervals of time based on recognizable geological and paleobiological markers, with typical length about 7 million years. As the figure shows,




Figure 1 Number of known marine families alive over the time interval from the Cambrian to the present. (The data are taken from Sepkoski, 1992)




Figure 2 Estimated extinction of marine animals in families per stratigraphic stage since the Cambrian. The arrows indicate the positions of the big five mass extinction events discussed in the text

there has been considerable variation in the intensity of extinction over prehistoric time. Of particular note are the five large peaks in extinction marked with arrows. These are the big five mass extinction events that marked the ends of the Ordovician, Devonian, Permian, Triassic, and Cretaceous periods. A sixth peak in the Cambrian is also visible, but this is thought probably to be an artifact of poor fossil preservation during that period rather than a real extinction event. The basic features of the big five extinctions are as follows (see also Table 1).

The end-Ordovician event about 440 million years ago appears to have occurred in two bursts, separated by about 1 million years, which between them wiped out about 85%

Table 1 Extinction intensities at the genus and species level for the big five mass extinctions of the Phanerozoic. Estimates of genus extinction are obtained from directed analysis of the fossil record while species loss is inferred using a statistical technique called reverse rarefaction. (Figures are taken from Jablonski, 1991)

Extinction	Genus loss (%) (observed)	Species loss (%) (estimated)
End Ordovician	60	85
Late Devonian	57	83
Late Permian	82	95
End Triassic	53	80
End Cretaceous	47	76

of then-living species. The event was confined to marine species, since multicellular life had not yet colonized the land. Particularly affected were brachiopods, bivalves, echinoderms, bryozoans, and corals. The immediate cause of extinction appears to have been the continental drift of a significant landmass into the south polar region, causing a global temperature drop, glaciation, and consequent lowering of the sea level, which destroyed species habitats around the continental shelves. The sea level rose again with the end of the glacial interval about a million years later and caused a second burst of extinction.

The late-Devonian extinction around 360 million years ago is complex and rather poorly understood. It is probably, in fact, composed of a number of separate events – as many as seven – spread over about 25 million years, including particularly notable extinctions at the ends of the Givetian, Frasnian, and Famennian stages. Overall, about 80% of living species died out in the late Devonian. Particularly hard hit were corals, brachiopods, bryozoans, ammonoids, and fish. The causes of these extinctions are unclear. The leading theories suggest that changes in sea level and ocean anoxia, possibly triggered by global cooling or oceanic volcanism, were most likely responsible, although the impact of an extraterrestrial body such as a comet has also been considered.

The late-Permian extinction around 250 million years ago was the largest extinction event of all time, killing some 95% of marine species and about 70% of land-dwelling species. Like the end-Ordovician event, it seems to have been composed of two bursts, separated by an interval of about 10 million years, the second being the larger of the two. Notable extinction happened again amongst brachiopods, ammonoids, and corals, as well as gastropods and, unusually, insects. Despite an enormous amount of research on the subject, the causes of the late-Permian event are still a subject of debate. It is clear, however, that the sea level rose during this period, levels of oxygen in the oceans were low, and carbon dioxide (CO₂) levels were high. There is some suggestion that a cometary impact may have been involved, or a shift in ocean circulation driven

by climate change, or CO₂ and sulfur release following large-scale volcanic activity. The late-Permian event had a profound effect on the terrestrial ecosystem, which is still being felt today, a quarter of a billion years later. A particularly notable example amongst marine faunas is that of the bivalves, a relatively minor group during the Paleozoic that took advantage of the ecological vacuum left by the extinction to establish a solid grip on shallow-water environments, leading to their dominance over the previously very successful brachiopods and gastropods.

The end-Triassic extinction around 210 million years ago is probably the most poorly understood of the big five extinction events. It appears to have killed about 80% of species then living, either in one burst or possibly in two, separated by about 20 million years. Major extinction is observed particularly amongst ammonoids, bivalves, gastropods, and brachiopods. Leading theories of the causes of the end-Triassic event are ocean anoxia, massive volcanism, or possibly an asteroid impact.

The end-Cretaceous event, usually called the *KT event*, has attracted the most popular interest of any extinction because it saw the death of those perennial movie stars, the dinosaurs, but it was in fact the smallest by quite a wide margin of all the big five. The *KT* event appears to have been a single pulse of extinction around 65 million years ago, which wiped out about 70% of all species then living. As well as the dinosaurs, it extinguished many other land-dwelling vertebrates, especially large-bodied ones, along with large numbers of (marine) bivalves, gastropods, and foraminifera. The proximal cause of the *KT* event was, almost certainly, the impact of a large comet or meteor near the present site of the town of Puerto Chicxulub on the Yucatán peninsula in eastern Mexico, with an associated drop in sea level and possibly short-term cooling or heating, or acid rain.

The fossil record can give us valuable insight into the nature of extinction and the effects of large-scale environmental change. It also indicates that recovery from extinction is a slow process by human standards, typically taking on the order of 5 or 10 million years. Thus, it is important to fend off such extinctions before they happen, rather than hoping that the ecosystem will prove robust enough to take care of itself.

Comparison between fossil and present-day extinction is not straightforward since, as mentioned above, the fossil record consists largely of marine organisms, whereas interest in contemporary extinction focuses mostly on land-dwelling organisms. Also recently extinct or currently endangered species tend to be rare, whereas the fossil record primarily reflects the most abundant and numerous biotas. Still, there are a number of general patterns in the fossil extinction record that may help us in the conservation of modern-day biodiversity.

First, we note that habitat loss, such as the destruction of shallow-water environments on the continental shelf as a result of changes in sea level, appears to have been an important cause of extinction again and again. A rise in sea level as a result of global warming over the next century, for example, could be devastating for reef communities. In this context, human-driven habitat loss is leading to extensive habitat fragmentation, i.e., to the generation of spatially isolated subareas. In many cases the subdivision of a large population into many weakly connected subpopulations substantially increases extinction risk. Again, the fossil record is very useful in providing a source of comparison. Habitat fragmentation enhanced the extinction of large mammals during the Pleistocene (75% of them are extinct). The area reduction during glacial cycles led to widespread habitat fragmentation and eventually to the extinction of many species. A considerable number of current endemisms in plant species are due to the confinement to small areas during the Ice Age.

Second, most past extinction events appear to have been selective to some extent. The end-Ordovician event, which was associated with a period of global cooling and glaciation, particularly favored species that were well adapted to cold-water conditions, and was particularly harsh on those who were not, for obvious reasons. The *KT* boundary event, as mentioned previously, appears to have come down especially hard on large-bodied animals. A number of explanations for this latter effect have been put forward. Large-bodied animals have smaller populations and greater area requirements, and are thus more sensitive to habitat loss or fragmentation. Moreover, their trophic requirements and low rates of population growth make them slow to recover from environmental change. It has been suggested that the loss of large herbivorous faunas could trigger major changes in biogeographic vegetation patterns that could in turn trigger further extinctions.

This last observation leads us to an additional question: to what extent do interactions between species affect the response of an ecosystem to environmental stress? Ecological interdependence between species may have heightened the impact of some of the mass extinctions; some studies, for example, have suggested that the collapse of marine food chains at the end-Cretaceous contributed to the *KT* extinction event. The *KT* event had a rapid effect on most biotas and a subsequent long-term effect perhaps related to a decrease in primary productivity. This seems particularly likely in the case of marine biotas, which show a marked dip in the rate of accumulation of carbonates, especially calcium carbonate, following the *KT* boundary, indicating a decrease in productivity. The resulting decrease in food supply would then produce extinctions at higher trophic levels. Similar mechanisms may also have been at work during the human-driven end-Pleistocene extinctions of mastodons and mammoths in North America, which were




Figure 3 An example of coextinction: the introduction of the myxoma virus to rabbit warrens in England triggered the extinction of the Large Blue butterfly (*Maculina arion*), an already endangered, endemic species. Myxomatosis caused a dramatic decline in rabbit populations and a consequent surge in the abundance of a species of grass on which the rabbits previously grazed (Grass-I). Another previously dominant species of grass (Grass-II) was unable to compete and declined substantially in its abundance, giving rise to a shortage of nesting material for certain variety of ant (*Mynnica*). As a result of a symbiotic dependence between the butterfly *Maculina* and this species of ant, the butterfly then became extinct

associated with widespread changes in vegetation patterns and the disappearance of many other species. One may well ask whether the current extinction of large herbivorous species in African ecosystems will result in similar concurrent extinctions. It is also believed that symbiotic and

parasitic species are particularly vulnerable to extinction, as a result of their dependence upon a partner or host. Many living species may, in fact, already be doomed to extinction, because of the loss of a partner in an essential symbiotic relationship (Figure 3). The situation with parasites may be even worse, since most parasites are specific to a single species of host, and extinction amongst parasites often goes unnoticed, even though their importance in maintaining diversity has been stressed by many studies. Since all known animals and plants have some parasitic load, extinctions of parasites must be widespread in evolutionary history. The eggs of what are believed to be ectoparasitic mites have been discovered on fossilized dinosaur remains, indicating that coevolutionary parasitism, and hence coextinction (the coupled extinction of ecological partners) is an ancient phenomenon. Estimating extinction rates for parasites is unfortunately difficult, since they are poorly represented or difficult to identify in the fossil record, and internal parasites are basically not preserved at all.

Overall, recent extinction rates in most plant and animal taxa are relatively low, but for some groups they do approach the levels associated with prehistoric mass extinction events (Table 2). These observations have led some researchers to suggest that the biosphere may be on the verge of another such event. Some claim that the world has already entered a sixth period of mass extinction, driven primarily by the human population explosion. Theoretical studies of long-term ecological responses to habitat destruction suggest that steady increases in extinction rates are to be expected in the near future. Paleontological studies indicate that rare, localized, or specialized species that have evolved to survive in particular niches are the mostly likely to become extinct, while widespread or adaptable species, or opportunistic colonizers are likely to prevail. The fossil record is thus more than a cautionary tale.

Table 2 Comparison between different features of past mass extinction events (as reported from the fossil record) and present-day, human-driven mass extinction. Here the effects of habitat loss are widespread but different mechanisms were at work: changes in sea levels and continental breakup (fossil record) and human-driven habitat destruction/fragmentation (present day) respectively. (Estimations of species loss are from Wilson, 1992)

	Fossil record	Present day
Time resolution	~10 ⁵ –10 ⁶ years	1 year
Most affected biotas	Tropical biotas	Coral reefs, rainforests
Selectivity on size	Large-sized species	Large-sized species
Loss of endemics	Not well known	Widespread
Effects of habitat loss	Widespread	Widespread
Time scale of recovery	~5–10 Myr	Not known
Direct effects on food webs	Very important	Very important
Indirect effects on food webs	Not known	Very important
Extinction rates	75–80%	~10 ⁴ species year ⁻¹

As paleontologist David Jablonski has put it:

The lessons from the past are inevitably blurry at a coarse scale. At the present stage of knowledge, the fossil record is more revealing for potential long-term consequences than for immediate solutions. However, the history of life of Earth provides an array of worst-case scenarios ... that are sufficiently spectacular to militate against inaction.

REFERENCES

- Jablonski, D (1991) Extinctions: A Paleontological Perspective, *Science*, **253**, 754–757.
- Sepkoski, Jr, J J (1992) *A Compendium of Fossil Marine Animal Families, Milwaukee Public Museum Contributions in Biology*

and Geology, **83**, 2nd edition, Milwaukee Public Museum, Milwaukee, WI.

- Wilson, E O (1992) *The Diversity of Life*, Harvard University Press, Cambridge, MA.

FURTHER READING

- Lawton, J H and May, R M, eds (1995) *Extinction Rates*, Oxford University Press, Oxford.
- Leakey, R and Lewin, R (1995) *The Sixth Extinction*, Doubleday, New York.
- Mckinney, M and Drake, J, eds (1998) *Biodiversity Dynamics*, Columbia University Press, New York.